

Table 6. Salmonella in laying hen flocks of Gallus gallus during the production period (flock-based data) in countries running control programmes, 2013

Country	Tested	Percent positive	S. Enteritidis S. Typhimurium % ^(a)	S. Enteritidis %	S. Typhimurium %	Other than SET %
Austria	2731	2.12	0.84	0.51	0.33	1.54
Belgium	606	5.94	1.82	1.65	0.17	4.62
Bulgaria	455	1.32	0	0	0	1.32
Croatia	322	2.8	2.8	2.8	0	0
Cyprus	40	87.5	7.5	7.5	0	80
Czech Republic	471	2.55	1.49	1.49	0	1.06
Denmark	373	1.07	1.07	0.27	0.8	0
Estonia	33	3.03	3.03	0	3.03	0
Finland	844	0	0	0	0	0
France	4974	0.6	0.6	0.34	0.26	0
Germany	5338	2	1.18	0.75	0.43	0.82
Greece	432	6.71	1.16	1.16	0	6.02
Hungary	1055	6.45	1.99	1.8	0.19	4.45
Ireland	208	0	0	0	0	0
Italy	2277	2.55	1.27	1.01	0.26	8.83
Latvia	44	2.27	2.27	2.27	0	0
Lithuania	97	0	0	0	0	0
Luxembourg	7	0	0	0	0	0
Malta	85	49.41	1.18	1.18	0	48.24
Netherlands	3457	0.72	0.72	0.55	0.17	0
Poland	2413	3.98	2.4	2.2	0.21	1.57
Portugal	383	6.27	1.57	1.57	0	5.22
Romania	4596	2.44	0.81	0.81	0	1.63
Slovakia	396	1.26	0.25	0.25	0	1.01
Slovenia	182	3.85	1.1	1.1	0	2.75
Spain	2135	8.76	1.87	1.55	0.33	6.89
Sweden	636	1.1	1.1	0	1.1	0
United Kingdom	4012	0.92	0.07	0.05	0.02	0.85
Iceland	30	0	0	0	0	0
Norway	747	0	0	0	0	0
Switzerland	901	0.22	0.2	0.11	0.11	0
Total (MS)	38602	2.5%	1	0.7%	0.22	2.0%

Target (production period) is calculated from the prevalence reported in 2012. Target is set up at 2.0 % for most of the countries, with the exception of the following: Cyprus (11.0 %), Malta (5.5 %), Luxembourg (3.2 %) and Poland (2.6 %). Croatia did not have the relative reduction target for 2013, as 2013 was the first year of reporting for this MS.

(a): S. Typhimurium includes monophasic S. Typhimurium.